

Complexes of N-thiophosphorylthiourea $\text{tBuNHC(S)NHP(S)(OiPr)}_2$ with zinc(II), cadmium(II), nickel(II), and cobalt(II) cations

Safin D., Sokolov F., Gimadiev T., Brusko V., Babashkina M., Chubukaeva D., Krivolapov D., Litvinov I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Reaction of the potassium salt of N-thiophosphorylthiourea $\text{tBuNHC(S)NHP(S)(OiPr)}_2$ (HL) with ZnII , CdII , NiII and CoII in aqueous EtOH leads to complexes of common formula M(L-S,S')_2 (ML_2). Complexes were investigated by IR, UV-Vis, ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy and microanalysis. The structure of complex NiL_2 was investigated by single crystal X-ray diffraction analysis. The nickel(II) ion has a square-planar environment, S_4 , with two anionic ligands involving 1,5-S,S'-coordination mode. The ligands are bound in a trans configuration. © 2008 Wiley-VCH Verlag GmbH & Co. KGaA.

<http://dx.doi.org/10.1002/zaac.200700549>

Keywords

Cadmium, Chelates, Cobalt, Crystal structures, N-thiophosphorylthiourea, Nickel, Zinc